

SINP/TNP/93-23

hep-th/9312128

STRUCTURE OF MULTI-ANYON WAVEFUNCTIONS

P. Mitra¹

Saha Institute of Nuclear Physics

Block AF, Bidhannagar

Calcutta- 700 064, INDIA

Abstract

Symmetries of multi-anyon wavefunctions are analysed with the help of a second quantized formulation. Analogues of Slater determinants are constructed. It is shown that the Pauli principle is not enforced.

¹e-mail address mitra@saha.ernet.in

The description of states containing two or more identical particles is a standard part of quantum mechanics books [1]. It has been well known for a long time now that wavefunctions of such states behave in one of two possible ways: they are either symmetric or antisymmetric under exchange of coordinates of the particles. The two cases correspond to Bose and Fermi statistics respectively. But if one considers a spacetime which is (2+1) dimensional, the situation becomes more complicated [2]. There is no sharp division into two classes here; one can have a continuous range of statistics, often called *anyon* statistics. Anyons, which exist only in (2+1) dimensions, interpolate between Bose and Fermi statistics.

Although they exist in a somewhat exotic spacetime, anyons are likely to be physically important, with significant condensed matter applications. That is why this subject has been keenly followed in the recent literature [3]. Quite a lot is known already about the quantum mechanics of anyons [4]. Some progress has also been made in the field theory of these objects [5]. However, the behaviour under exchange of particle coordinates of multianyon wavefunctions has not been properly studied in the light of the field theoretic formulation. This is what is proposed to be done in this note.

When a particle is taken round an identical particle, it picks up a factor which is ± 1 for bosons and fermions, and some phase factor for anyons. It follows that instead of having a wavefunction that is symmetric or antisymmetric under the exchange of coordinates, two or more anyons can only have a wave-

function which changes by a phase under such an exchange. The modulus of the wavefunction must remain unaltered. But the modulus is not a linear function of the wavefunction, so this statement is not particularly useful, and a more detailed description is necessary. The phase factor associated with an exchange of identical anyons may be taken to be $q = e^{i\theta}$. The parameter q or θ will of course depend on the type of anyon considered. For instance, in the limiting cases of bosons and fermions, θ can be taken to be 0 or 1. But there is an extra complication for anyons. In general, the inverse of $q = e^{i\theta}$ is unequal to q , so that the exchange has to be assigned a direction or sense. Correspondingly, the transformation group that enters the discussion is not the permutation group but the *braid* group. We shall have more to say about these groups later. Let us note now that in the literature, the dependence of the phase factor on the direction of the exchange is taken into account by specifying whether the rotation is clockwise or anticlockwise. This corresponds to the use of multivalued wavefunctions. We prefer an alternative description where a *cut* is chosen and wavefunctions become single-valued but discontinuous. Then, instead of writing simply q , it is necessary to introduce $q(\vec{x}, \vec{y})$, which is equal to q or q^{-1} depending on the relative positions $\vec{x}, \vec{y}$.

This is best understood by referring to the second quantized picture and writing

$$a^\dagger(\vec{x})a^\dagger(\vec{y}) = q(\vec{x}, \vec{y})a^\dagger(\vec{y})a^\dagger(\vec{x}), \quad (1)$$

where creation operators have been taken to exist. It is believed that there has

to be some amount of nonlocality in theories with anyons. But this nonlocality does not manifest itself in simple commutation relations like this, which are familiar in model field theories containing anyonic excitations. Self-consistency demands that

$$q(\vec{x}, \vec{y}) = q(\vec{y}, \vec{x})^{-1}. \quad (2)$$

In the usual situation of Bose or Fermi statistics, where $q^2 = 1$, q is independent of location and therefore continuous. But in the case of anyons with $q^2 \neq 1$, q is a discontinuous function of its arguments. One possibility that can be visualized for the cut is such that

$$q(\vec{x}, \vec{y}) = e^{i\theta\epsilon(\alpha x_1 + \beta x_2 - \alpha y_1 - \beta y_2)}, \quad (3)$$

where ϵ represents the sign function. This is self-consistent but discontinuous.

We shall not need the explicit form of $q(\vec{x}, \vec{y})$.

In general, by translation invariance, $q(\vec{x}, \vec{y})$ can depend only on the difference $\vec{x} - \vec{y}$. Further, eq. (2) leads to

$$q(\vec{x}, \vec{x}) = \pm 1. \quad (4)$$

These two possibilities can be referred to as the cases of *bosonic* anyons and *fermionic* anyons respectively, if there is no confusion. In the literature, *fermionic* anyons are predominant. It should be emphasized that this distinction and this nomenclature refer exclusively to the behaviour at zero separation and are not of fundamental importance.

We come now to the behaviour of 2-anyon wavefunctions under the exchange of particle coordinates. A wavefunction $\psi(\vec{x}, \vec{y})$ corresponds to the 2-anyon creation operator

$$\int d\vec{x} \int d\vec{y} \psi(\vec{x}, \vec{y}) a^\dagger(\vec{x}) a^\dagger(\vec{y}). \quad (5)$$

Now ψ can be decomposed uniquely into what may be called its *q-symmetric* and *q-antisymmetric* parts:

$$\begin{aligned} \psi(\vec{x}, \vec{y}) &= \psi_{q \text{ sym}}(\vec{x}, \vec{y}) + \psi_{q \text{ ant}}(\vec{x}, \vec{y}) \\ &\equiv \frac{1}{2}[\psi(\vec{x}, \vec{y}) + q(\vec{y}, \vec{x})\psi(\vec{y}, \vec{x})] + \frac{1}{2}[\psi(\vec{x}, \vec{y}) - q(\vec{y}, \vec{x})\psi(\vec{y}, \vec{x})]. \end{aligned} \quad (6)$$

The above creation operator can therefore be written as

$$\int d\vec{x} \int d\vec{y} (\psi_{q \text{ sym}}(\vec{x}, \vec{y}) + \psi_{q \text{ ant}}(\vec{x}, \vec{y})) a^\dagger(\vec{x}) a^\dagger(\vec{y}). \quad (7)$$

Now by interchanging the integration variables $\vec{x}$ and $\vec{y}$, and making use of eq. (1), one sees that

$$\begin{aligned} &\int d\vec{x} \int d\vec{y} \psi_{q \text{ ant}}(\vec{x}, \vec{y}) a^\dagger(\vec{x}) a^\dagger(\vec{y}) \\ &= \int d\vec{x} \int d\vec{y} \frac{1}{2} \psi(\vec{x}, \vec{y}) [a^\dagger(\vec{x}) a^\dagger(\vec{y}) - q(\vec{x}, \vec{y}) a^\dagger(\vec{y}) a^\dagger(\vec{x})] \\ &= 0. \end{aligned} \quad (8)$$

This shows that the *q-antisymmetric* part of ψ does not contribute to this operator, so that ψ can be taken to be *q-symmetric* without loss of generality. Thus the generalization of the symmetry or antisymmetry of two-particle wavefunctions to the anyonic case is the *q-symmetry* introduced in eq. (6).

The reader should be warned to be careful with the order of the arguments in these definitions. With the convention chosen above, the product $a^\dagger(\vec{x})a^\dagger(\vec{y})$ is *not* q -symmetric. Indeed, eq. (6) shows that

$$\psi_{q \text{ sym}}(\vec{x}, \vec{y}) = q(\vec{y}, \vec{x})\psi_{q \text{ sym}}(\vec{y}, \vec{x}), \quad (9)$$

whereas eq. (1) has $q(\vec{x}, \vec{y})$ in it. Since the interchange of the two arguments of q amounts to inverting it, one concludes that $a^\dagger(\vec{x})a^\dagger(\vec{y})$ is in fact q^{-1} -symmetric. The lesson is that *if a q^{-1} -symmetric object is contracted with anything, that object can be taken to be q -symmetric without loss of generality.* For $q = 1$, this is of course a well-known result.

The generalization to states of more than two particles is not difficult, but there are some extra complications. To study 3-anyon states, note that the product $a^\dagger(\vec{x})a^\dagger(\vec{y})a^\dagger(\vec{z})$ is q^{-1} -symmetric in $\vec{x}$ and $\vec{y}$, and also in $\vec{y}$ and $\vec{z}$, but there is no simple symmetry property corresponding to the interchange of $\vec{x}$ and $\vec{z}$. However, this interchange can be written in terms of the other two interchanges, so that a formula can be constructed to express the symmetry property corresponding to this interchange. The real question is about the wavefunction. By virtue of the above proposition regarding q -symmetric objects, it is clear that the wavefunction $\psi(\vec{x}, \vec{y}, \vec{z})$ is q -symmetric in its first two arguments and also in its last two arguments:

$$\begin{aligned} \psi(\vec{x}, \vec{y}, \vec{z}) &= q(\vec{y}, \vec{x})\psi(\vec{y}, \vec{x}, \vec{z}) \\ &= q(\vec{z}, \vec{y})\psi(\vec{x}, \vec{z}, \vec{y}). \end{aligned} \quad (10)$$

From this it follows that

$$\psi(\vec{x}, \vec{y}, \vec{z}) = q(\vec{y}, \vec{x})q(\vec{z}, \vec{x})q(\vec{z}, \vec{y})\psi(\vec{z}, \vec{y}, \vec{x}). \quad (11)$$

Although the exchange of the first and the third arguments does amount to multiplying the wavefunction by a phase factor, *this phase factor depends on the second argument too*. This kind of relation can also be found for wavefunctions of states with even more anyons.

These symmetries can be used to construct many -anyon wavefunctions starting from those for single anyons. If $\psi_1(\vec{x})$ and $\psi_2(\vec{x})$ are two such wavefunctions, a "noninteracting" 2-anyon wavefunction may in principle be found by multiplying these functions and appropriately *q-symmetrizing* the product. It is assumed here that a Hamiltonian constructed by adding two single -anyon Hamiltonians without any interaction gives a reasonable description of reality. From eq. (6), the wavefunction is found to be

$$\frac{1}{2}[\psi_1(\vec{x})\psi_2(\vec{y}) + q(\vec{y}, \vec{x})\psi_1(\vec{y})\psi_2(\vec{x})]. \quad (12)$$

This expression has two terms corresponding to the two ways of permuting the particles. However, unlike the usual cases of Bose or Fermi statistics, the superposition involves the nonconstant function q . For $q = \pm 1$, the usual symmetrized and antisymmetrized expressions are of course obtained. Since for $q = -1$ a determinant appears, one can, in the general case, speak of a *(-q) determinant* (apart from a factor of a half).

The 3-anyon wavefunction is more complicated. In terms of three single

-anyon wavefunctions, one can construct the appropriately q -symmetrized product

$$\begin{aligned}
& \frac{1}{3!} [\psi_1(\vec{x})\psi_2(\vec{y})\psi_3(\vec{z}) + q(\vec{y}, \vec{x})\psi_1(\vec{y})\psi_2(\vec{x})\psi_3(\vec{z}) + q(\vec{z}, \vec{y})\psi_1(\vec{x})\psi_2(\vec{z})\psi_3(\vec{y}) \\
& + q(\vec{z}, \vec{y})q(\vec{z}, \vec{x})\psi_1(\vec{z})\psi_2(\vec{x})\psi_3(\vec{y}) + q(\vec{y}, \vec{x})q(\vec{z}, \vec{x})\psi_1(\vec{y})\psi_2(\vec{z})\psi_3(\vec{x}) \\
& + q(\vec{y}, \vec{x})q(\vec{z}, \vec{x})q(\vec{z}, \vec{y})\psi_1(\vec{z})\psi_2(\vec{y})\psi_3(\vec{x})]
\end{aligned} \tag{13}$$

There are six terms here corresponding to the six ways of permuting the particles, and once again there are nontrivial location -dependent coefficients for these terms. When $q = 1$, the product is simply the symmetrized product, and for $q = -1$, the usual determinant appears. In general one can refer to the expression in the square brackets as a $(-q)$ *determinant*. From these two examples, the construction of $(-q)$ *determinants* for an arbitrary number of identical anyons should be clear.

A comment about the Pauli principle is in order. In the case of fermions, the fact that $q = -1$ leads to the vanishing of a 2 -particle wavefunction for coincident arguments,

$$\begin{aligned}
\psi(\vec{x}, \vec{x}) &= -\psi(\vec{x}, \vec{x}) \\
&= 0,
\end{aligned} \tag{14}$$

and this argument is thought to generalize to the case of anyons where q is not in general equal to unity. However, as argued above in eq. (4), the *relevant* value of q is ± 1 depending on whether the anyons are of the *bosonic* or the *fermionic* type, so this generalization holds only for the latter type. There is no

need for the wavefunction of two *bosonic* anyons to vanish when the coordinates coincide. More important, the *q-symmetrized* products constructed above are in general nonvanishing even when two or more of the single-anyon wavefunctions coincide, and this is true for both *fermionic* and *bosonic* anyons. This proves that the Pauli exclusion principle does *not* hold in the general case of anyons and is restricted to the case of fermions for which it was after all intended.

Lastly, some comments have to be made about the supersession of the permutation group by the braid group in the case of anyons. For usual statistics, a multiparticle wavefunction changes by ± 1 under an interchange of particles, the sign factor being related to the statistics and the permutation involved. For bosons, all permutations give rise to the factor $+1$, while for fermions, the sign factor is $(-1)^P$, where P is the parity (even or odd) of the permutation. In both cases, the permutation group is represented one-dimensionally on the multiparticle wavefunction. In the case of anyons, the factors which appear when two particles are interchanged are phase factors like q and their products. In general, these phase factors do *not* form a group. To have a group structure, squares and higher powers of such phase factors would have to be accommodated. Such a situation could arise if multivalued wavefunctions were used. But we have preferred to introduce cuts and stick to (discontinuous) single-valued wavefunctions. A representation of the braid group is *not* obtained.

It is a pleasure to thank Avinash Khare for asking a question and thereby initiating this analysis.

References

- [1] A. Messiah, *Quantum Mechanics*, North- Holland, Amsterdam (1961)
- [2] J. Leinaas and J. Myrheim, Nuov. Cim. **B37**, 1 (1977); G. Goldin, R. Menikoff and D. Sharp, Journ. Math. Phys. **22**, 1664 (1981); F. Wilczek, Phys. Rev. Lett. **48**, 1144 (1982)
- [3] S. Forte, Rev. Mod. Phys. **64**, 193 (1992)
- [4] A. Khare and J. McCabe, Phys. Lett. **B269**, 330, (1991); M. Sporre, J. Verbaarschot and I. Zahed, Phys. Rev. Lett. **67**, 1813, (1991); M. V. N. Murthy, J. Law, M. Brack and R. Bhaduri, Phys. Rev. Lett. **67**, 1817, (1991); S. A. Chin and C. R. Hu, Phys. Rev. Lett. **69**, 229 (1992) and **69**, 1148E (1992)
- [5] G. Semenoff, Phys. Rev. Lett. **61**, 517 (1988); J. Fröhlich and P. Marchetti, Comm. Math. Phys. **121**, 177 (1989); M. Lüscher, Nucl. Phys. **B326**, 557 (1989); D. Eliezer, G. Semenoff and S. S. C. Wu, Mod. Phys. Lett. **A7**, 513 (1992); R. Banerjee, A. Chatterjee and V. Sreedhar, Ann. Phys., **222**, 254 (1993); P. Mitra, Phys. Lett. **B313**, 41 (1993)